For Clinics:

How to identify and diagnose VCF's

Patient Presents in Clinic

- Complain of sudden onset of back pain
- Patients at risk include those with osteoporosis, long term steroid use, known or occult malignancies



Physical Exam

- Usually involve localized pain
- Start with light palpation of the spinous process and proceed to percussion as indicated
- Focused neurological examination recommended to exclude acute neural injury to the spinal cord or nerve roots



If VCF is Suspected...

- Recommend a lateral plain film x-ray
- MRI to confirm & r/o retropulsed fragment
- Bonescan with spect if cannot do MRI
- May consider CT



If VCF is present...

- Localize pain & administer appropriate pain medication
- Contact Pain Care Boise



Dr. William G. Binegar, MD



Welcome Dr. Vaughn!

Dr. Vaughn completed his anesthesiology and interventional pain management fellowship at the Texas Tech University Health Sciences Center. He is a member of the of the American Board of Anesthesiology as well as the American Society of Interventional Pain.



(208) 342-8200

301 W Myrtle St, Boise, ID 83702 Idaho

www.paincareboise.com



Pain Care BOISE

Medicare Certified

AAAHC Certified

Risks of Conservative Treatment for VCF

(Vertebral Compression Fractures)

The "Who" of the Problem


- 50% of women and 25% of men will have an osteoporotic fracture in their lifetime¹
- 10 million Americans have osteoporosis²
- Additional 34 million have osteopenia²
- 700,000 osteoporotic fractures occur annually in the U.S.³



VCF (Vertebral Compression Fractures) when it goes untreated

Risks of Conservative Therapy

- Bed rest
 - Leading to "the downward spiral"
- Bracing
- Open surgery fixation equipment
 Pain tolerance



What's new:

Medicare No Longer Requires 6 Weeks of Conservative Therapy for VCFs

- Medicare Administrative Contractors (MAC) CGS, NGS, Noridian and WPS have changed the VCF LCD that will alter Medicare vertebroplasty and vertebral augmentation coverage beginning in mid to late 2020.
- Coverage will no longer require 6
 weeks of conservative therapy and
 will now cover acute fractures (< 12
 weeks) with no conservative
 management required.
 - Ballon Kyphoplasty patients experience lower Mortality rate and overall morbididty than conservative management VCF patients.

Studies conclude that for every 15 patients treated with VCF, one life is saved. (Edidin et all)

Reduce Mortality from VCF's by contacting Pain Care Boise at (208) 342-8200

https://www.nof.org/patients/what-is-osteoporosis/(2017).
 May 11, 2015.

 [&]quot;10 Facts About Osteoporosis and Bone Health." Fact About Osteoporosis. ShareCare, 1 Jan. 2015. Web.
 Siemionow K, Lieberman IH. Vertebral Augmentation in Osteoporotic and Osteolytic Fractures. Curr Opin. Support Palliat Care. 2009 Sep;3(3):219-25